

DP.C.2. Use of species and severity

	Non-recovery	%	Mild [up to and including]	%	Moderate	%	Severe	%	Total
Mice	14,205	8.63%	91,615	55.69%	57,566	34.99%	1,136	0.69%	164,522
Rats	6,015	15.62%	15,073	39.15%	17,177	44.62%	233	0.61%	38,498
Guinea-Pigs	317	10.76%	2,368	80.35%	260	8.82%	2	0.07%	2,947
Hamsters (Syrian)			3	1.06%	269	95.39%	10	3.55%	282
Hamsters (Chinese)									
Mongolian gerbil									
Other Rodents			3	100.00%					3
Rabbits	165	6.75%	520	21.29%	1,749	71.59%	9	0.37%	2,443
Cats									
Dogs	8	3.74%	152	71.03%	54	25.23%			214
Ferrets			4	100.00%					4
Other carnivores			229	22.13%	773	74.69%	33	3.19%	1,035
Horses, donkeys & cross-breeds			119	100.00%					119
Pigs	1,713	28.60%	2,290	38.24%	1,967	32.84%	19	0.32%	5,989
Goats					1	100.00%			1
Sheep	16	24.24%	40	60.61%	10	15.15%			66
Cattle	48	1.30%	3,392	91.82%	254	6.88%			3,694
Prosimians									
Marmoset and tamarins									
Cynomolgus monkey									
Rhesus monkey									
Vervets Chlorocebus spp.									
Baboons									
Squirrel monkey									
Other species of New World Monkeys (Ceboidea)									
Other species of Old World Monkeys (Cercopithecoidea)									
Other species of non-human primates									
Apes									
Other Mammals			28	100.00%					28
Domestic fowl			172	36.91%	294	63.09%			466
Other birds			369	90.44%	39	9.56%			408
Reptiles	139	48.10%	52	17.99%	98	33.91%			289
Rana			4	100.00%					4
Xenopus			60	21.05%	225	78.95%			285
Other Amphibians			47	100.00%					47
Zebra fish			1,467	92.44%			120	7.56%	1,587
Other Fish	245	1.57%	9,577	61.37%	5,584	35.78%	200	1.28%	15,606
Cephalopods									
Total	22,871		127,584		86,320		1,762		238,537

DP.C.2. Use of species and severity

